



## Role of clothing in both accelerating and impeding dermal absorption of airborne SVOCs

**Morrison, Glenn C.; Weschler, Charles J.; Bekö, Gabriel; Koch, Holger M.; Salthammer, Tunga; Schripp, Tobias; Toftum, Jørn; Clausen, Geo**

*Published in:*  
Journal of Exposure Science and Environmental Epidemiology

*Link to article, DOI:*  
[10.1038/jes.2015.42](https://doi.org/10.1038/jes.2015.42)

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Morrison, G. C., Weschler, C. J., Bekö, G., Koch, H. M., Salthammer, T., Schripp, T., Toftum, J., & Clausen, G. (2016). Role of clothing in both accelerating and impeding dermal absorption of airborne SVOCs. *Journal of Exposure Science and Environmental Epidemiology*, 26(1), 113-118. <https://doi.org/10.1038/jes.2015.42>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Title:** Role of clothing in both accelerating and impeding dermal absorption of airborne SVOCs

**Authors:** Glenn C. Morrison, PhD<sup>1</sup>, Charles J. Weschler, PhD<sup>2,3</sup>, Gabriel Bekö, PhD<sup>3</sup>, Holger M. Koch, PhD<sup>4</sup>, Tunga Salthammer, PhD<sup>5</sup>, Tobias Schripp, PhD<sup>5</sup>, Jørn Toftum, PhD<sup>3</sup> and Geo Clausen, PhD<sup>3</sup>

**Affiliations:**

<sup>1</sup> Civil, Architectural and Environmental Engineering, Missouri University of Science and Technology, Rolla, MO, 65409, USA

<sup>2</sup> Environmental and Occupational Health Sciences Institute, Rutgers University, 170 Frelinghuysen Road, Piscataway, NJ 08854, USA

<sup>3</sup> International Centre for Indoor Environment and Energy, Department of Civil Engineering, Technical University of Denmark, DK-2800, Lyngby, Denmark

<sup>4</sup> Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr-Universität Bochum (IPA), Bürkle-de-la-Camp-Platz 1, 44789 Bochum, Germany

<sup>5</sup> Fraunhofer WKI, Department of Material Analysis and Indoor Chemistry, Bienroder Weg 54E, 38108 Braunschweig, Germany

**Address correspondence to:** Glenn Morrison, Civil, Architectural and Environmental Engineering, Missouri University of Science and Technology, Rolla, MO, USA.

Telephone: 573-341-7192. Fax: 573-341-4729. E-mail: [gcm@mst.edu](mailto:gcm@mst.edu).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

24  
  
25  
  
26  
  
27  
  
28  
  
29  
  
30

**Running title:** Clothing enhances dermal uptake of airborne SVOCs

**Competing financial interests:** All authors declare no actual or potential competing financial interests.

For Peer Review Only

31

**Abstract**

To assess the influence of clothing on dermal uptake of SVOCs, we measured uptake of selected airborne phthalates for an individual wearing clean clothes or air-exposed clothes and compared these results with dermal uptake for bare-skinned individuals under otherwise identical experimental conditions. Using a breathing hood to isolate dermal from inhalation uptake, we measured urinary metabolites of diethylphthalate (DEP) and di-n-butylphthalate (DnBP) from an individual exposed to known concentrations of these compounds for 6 hours in an experimental chamber. The individual wore either clean (fresh) cotton clothes or cotton clothes that had been exposed to the same chamber air concentrations for 9 days. For a 6-hour exposure, the net amounts of DEP and DnBP absorbed when wearing fresh clothes were respectively 0.017 and 0.007  $\mu\text{g}/\text{kg}/(\mu\text{g}/\text{m}^3)$ ; for exposed clothes the results were 0.178 and 0.261  $\mu\text{g}/\text{kg}/(\mu\text{g}/\text{m}^3)$  (values normalized by air concentration and body mass). When compared against the average results for bare-skinned participants, clean clothes were protective, while exposed clothes increased dermal uptake for DEP and DnBP by factors of 3.3 and 6.5 respectively. Even for non-occupational environments, wearing clothing that has adsorbed/absorbed indoor air pollutants can increase dermal uptake of SVOCs by substantial amounts relative to bare skin.

**Introduction**

Dermal absorption of organic compounds directly from air has been observed for some volatile and semi-volatile compounds. In reviews by Rehal et al.<sup>1</sup> and Rauma et al.<sup>2</sup> a handful of volatile organic compounds (VOCs) have been observed to have dermal uptakes

1  
2  
3 54 that are substantial compared to inhalation intakes. For example, Piotrowski<sup>3,4</sup> found that  
4  
5 55 nitrobenzene and phenol doses via dermal absorption were about 50% those due to  
6  
7  
8 56 absorption by inhalation. Weschler and Nazaroff<sup>5,6</sup> argued that the dermal absorption dose  
9  
10 57 from air could also compare with or exceed the dose due to inhalation for semi-volatile  
11  
12  
13 58 organic compounds (SVOCs) that meet specific criteria under steady-state conditions. In a  
14  
15 59 refinement of that model for non-steady-state conditions, Gong et al.<sup>7</sup> showed that timing of  
16  
17 60 exposure can significantly influence dose due to resistance and accumulation within the  
18  
19  
20 61 dermis. In a test of the hypothesis that the dermal dose of SVOCs from air could be  
21  
22  
23 62 significant, Weschler et al.<sup>8</sup> showed that dermal absorption was approximately equal to  
24  
25 63 inhalation dose for six bare-skinned male participants exposed to diethylphthalate (DEP)  
26  
27 64 and di n-butylphthalate (DnBP) for six hours in a chamber.  
28  
29  
30 65

31  
32 66 A few studies have evaluated how clothing may influence dermal uptake of organic  
33  
34 67 compounds from air or by transfer from treated fabrics. Piotrowski<sup>3</sup> found that clothing  
35  
36 68 reduced dermal uptake of airborne nitrobenzene by about 20-30% but had no observable  
37  
38  
39 69 effect on phenol absorption.<sup>4</sup> Organics that have been applied to clothing can be also  
40  
41  
42 70 absorbed. Blum et al.<sup>9</sup> observed metabolites of a flame retardant in the urine of children  
43  
44 71 who had worn clothing treated with this flame retardant. Similarly, subjects wearing  
45  
46 72 permethrin-impregnated battle dress uniforms absorbed this insecticide as evidenced by  
47  
48  
49 73 urinary metabolites<sup>10-12</sup>.

50  
51 74  
52  
53  
54 75 We hypothesize that sorption to clothing acts either to reduce or to increase dermal  
55  
56 76 uptake, depending on the extent to which the clothing has equilibrated with room air  
57  
58  
59  
60

1  
2  
3 77 contaminants prior to wearing. For some compounds, the boundary layer of air adjacent to  
4  
5  
6 78 the skin presents greater resistance to transport than does the stratum corneum and viable  
7  
8 79 epidermis.<sup>6</sup> For such compounds uptake is sensitive to the magnitude of the boundary layer  
9  
10  
11 80 permeability<sup>7</sup> and could be altered significantly by sorption to fabrics, especially for  
12  
13 81 compounds with high air-fabric partition coefficients.  
14  
15  
16 82

17  
18 83 For fabrics that are initially clean, adsorption to fabric fibers should decrease fabric  
19  
20 84 permeability, and lower overall dermal uptake, by reducing diffusional flux through the  
21  
22 85 fabric. With continued exposure, fabric permeability would increase as fabric surfaces  
23  
24 86 equilibrate with SVOCs. Fabrics that are exposed to building air for extended periods (e.g.  
25  
26 87 hanging up in a closet) may absorb a substantial quantity of SVOCs, or even reach  
27  
28 88 equilibrium, prior to wearing. For these clothes, we predict that dermal uptake will be  
29  
30 89 higher than uptake to bare skin.  
31  
32  
33  
34  
35 90

36  
37 91 Our objective is to test this hypothesis with two compounds that have been predicted, and  
38  
39 92 recently shown, to exhibit low dermal uptake resistance relative to mass-transfer  
40  
41 93 resistance through the layer of air adjacent to body surfaces. In this study, we measure  
42  
43 94 urinary concentrations and total excretion of DEP and DnBP metabolites during and after a  
44  
45 95 participant is exposed for 6 hours to known air concentrations of DEP and DnBP for 2  
46  
47 96 conditions: i) wearing freshly cleaned cotton clothing; ii) wearing previously clean cotton  
48  
49 97 clothing that had been exposed to the phthalates for at least one week. Inhalation uptake is  
50  
51  
52 98 controlled with a breathing hood. Results are compared against results from six individuals  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 99 who wore only shorts but were subjected to nearly identical conditions (results reported in  
4  
5  
6 100 Weschler et al.<sup>8</sup>).  
7

8 101

9  
10 102 **Methods**  
11

12  
13 103 The experiments reported here were integrated into the dermal uptake experiments  
14  
15 104 reported by Weschler et al.<sup>8</sup> and nearly all procedures, conditions and analytical methods  
16  
17 105 are therefore identical. The clothed individual was exposed to phthalates in the same  
18  
19  
20 106 chamber at the same time as bare-skinned participants during two of the chamber  
21  
22 107 exposure intervals, specifically Wednesday of the 1<sup>st</sup> week and Tuesday of the 2<sup>nd</sup> week.  
23  
24

25 108

26  
27 109 *Exposure chamber*  
28

29  
30 110 The 55 m<sup>3</sup> chamber housed two mixing fans, desks and chairs. The air exchange rate was  
31  
32 111 maintained at 0.7 1/h and the temperature was controlled at 30°C. The relative humidity  
33  
34 112 was not controlled and ranged from 20 to 35% during the experiments. A breathing hood  
35  
36 113 (Amron International, Vista, CA, #8890 Oxygen Treatment Hood) was used so that the  
37  
38  
39 114 participant in the clothing experiments could breathe clean air from outside the chamber,  
40  
41 115 thus allowing for the separation of dermal from inhalation dose. See Figure S.1 for an image  
42  
43 116 of the participant wearing test clothing and the hood while seated in the experimental  
44  
45 117 chamber. Air concentrations of DEP and DnBP were maintained by continuous emission  
46  
47 118 from aluminum panels (total area of 12 m<sup>2</sup>) coated with Latex paint. The paint had been  
48  
49  
50 119 formulated with 1% DEP and 10% DnBP (by weight), and was used to deliver these  
51  
52  
53 120 phthalates into chamber air at a relatively constant emission rate.<sup>8,13</sup>  
54  
55

56 121  
57  
58  
59  
60

### 122 *Clothing and clothing preparation*

123 Clothing was purchased from two different clothing stores in Rolla, Missouri, USA. Each set  
124 included a cotton undershirt, a pair of cotton jeans, a long-sleeved cotton tee shirt, cotton  
125 underwear and cotton socks. Details such as size, style and manufacturer can be found in  
126 Table S.1.

127  
128 Two sets of clothing were prepared by washing all pieces at the same time in a standard  
129 clothes washer using unscented detergent. They were then dried in an electric dryer on the  
130 “medium” setting and each set was packaged separately in two layers of clean aluminum  
131 foil until use. During the first 6-hour exposure period, one set was worn directly from its  
132 package and is denoted “fresh”. Another set of clothing was exposed to chamber air for 9  
133 days and denoted as “exposed”. This exposure took place in the same chamber, under the  
134 same conditions and at the same time as bare-skin dermal uptake experiments occurred;  
135 the latter are described in Weschler et al.<sup>8</sup> The clothing was hung inside-out in the path of  
136 fans to improve transfer of phthalates from air to the clothing. The air concentration was  
137 measured during days 2 and 3 of the 9 day clothing-exposure interval. During these  
138 periods, the average concentrations for DEP were 250 and 233  $\mu\text{g}/\text{m}^3$  and that of DnBP  
139 was 123 and 114  $\mu\text{g}/\text{m}^3$ .

### 141 *Preparation of participant*

142 Because there were a limited number of breathing hoods available in the exposure  
143 chamber, it was only possible to study one clothed participant. The participant was a 48  
144 year old Caucasian male, 192 cm tall weighing 91 kg. The participant followed the same



1  
2  
3 145 restricted diet and restricted use of personal care products protocol described in Weschler  
4  
5  
6 146 et al.<sup>8</sup> These restrictions were intended to reduce background metabolites of DEP and  
7  
8 147 DnBP in the participants' urine. In brief, for 12 hours prior to exposure and 54 hours after  
9  
10 148 exposure began, the participant only ate Swedish dried bread and ate thick-rinded fruit  
11  
12 149 such as oranges, bananas and melons. He drank only tap water or tea made from tap water.  
13  
14 150 The participant showered without soaps or detergents 24 hours prior to the experiment  
15  
16 151 and showered without soaps again 48 hours after the beginning of an exposure. The  
17  
18 152 research protocol was approved by the Capital Region of Denmark Committee for Research  
19  
20 153 Ethics. The participant provided informed consent before participation and consented to  
21  
22 154 publication of his photo.  
23  
24  
25  
26  
27  
28  
29

30 156 *Description of exposure periods*

31  
32 157 The participant participated in two exposure experiments. The first took place on a  
33  
34 158 Wednesday coincident with the 2<sup>nd</sup> set of exposure experiments during the first week  
35  
36 159 described in Weschler et al.<sup>8</sup> The participant collected two urine samples on the morning of  
37  
38 160 the experiment. Immediately before entering the chamber the participant collected a urine  
39  
40 161 sample, changed into the "fresh" set of experimental clothes, donned a breathing hood and  
41  
42 162 entered the chamber at 11:00. The participant sat at a desk for most of the 6-hour exposure  
43  
44 163 period and left the chamber once briefly to collect a urine sample. At 17:00, the participant  
45  
46 164 left the chamber and changed into his normal clothing. Following this, the participant  
47  
48 165 maintained the restricted diet and personal product restrictions and collected all urine for  
49  
50 166 48 hours. The second exposure experiment took place on a Tuesday coincident with the 3<sup>rd</sup>  
51  
52 167 set of exposure experiments during the second week described in Weschler et al.<sup>8</sup> The  
53  
54  
55  
56  
57  
58  
59  
60

procedure was identical to the first experiment except that the participant changed into the “exposed” set of clothes before entering the chamber at 10:00 (leaving at 16:00).

#### *Analysis of air and urine*

Air concentrations of phthalates were determined by first collecting 6 L samples of air with Tenax-TA filled thermal desorption tubes, and analyzing by thermal desorption followed by gas chromatography using a mass selective detector. Phthalates were quantified using original standards. The concentrations in air and other conditions are tabulated in Weschler et al.<sup>8</sup>

Urine samples were weighed on the day of collection and stored in a freezer until they were shipped overnight to the Institute for Prevention and Occupational Medicine of the German Social Accident Insurance in Bochum, Germany. Urine samples were analyzed for mono-ethyl phthalate (MEP), a metabolite of DEP, as well as mono-n-butyl phthalate (MnBP) and 3OH-mono-n-butyl phthalate (3OH-MnBP), metabolites of DnBP. The concentrations of these metabolites were determined by two-dimensional high performance liquid chromatography coupled to tandem mass spectrometry (LC/LC-MS/MS) using internal isotope-labeled standards after enzymatic deconjugation of the phthalate metabolites from the glucuronidated form following methods published by Koch et al.<sup>14,15</sup> Other details of analytical methods can be found in Weschler et al.<sup>8</sup>

#### *Calculations*

1  
2  
3 190 Calculated total uptake of DEP or DnBP during the 6 hour exposure period was based on  
4  
5  
6 191 methods described by Koch et al.<sup>15-17</sup> and outlined in Weschler et al.<sup>8</sup> Metabolite  
7  
8 192 concentrations were converted to mass excreted and then converted to parent molecule  
9  
10 193 uptake using predetermined metabolic conversion factors. In the “fresh clothes”  
11  
12  
13 194 experiment there was very low overall dermal uptake of DEP and DnBP (see Results and  
14  
15 195 Discussion). To better quantify uncertainty in this case, background uptake has been  
16  
17 196 determined in a somewhat different manner than in Weschler et al.<sup>8</sup> For the present  
18  
19  
20 197 participant, little residual uptake from the 6-hour experiment remained, relative to  
21  
22  
23 198 background uptake, for the last four urinations of the fresh clothes experiment (collected  
24  
25 199 from 40.5 to 50.0 hours after exiting the chamber). Therefore, the average dose rate (total  
26  
27 200 dose/elapsed time) from these samples was subtracted from the dose rate calculated for  
28  
29  
30 201 each post-exposure sample for both fresh and exposed clothes experiments. This was  
31  
32 202 multiplied by the sample time interval, and the result from each interval summed, to  
33  
34  
35 203 determine the background-corrected total dose. Dermal uptake was also corrected for DEP  
36  
37 204 and DnBP measured in the breathing hood air (40.7 µg/m<sup>3</sup> and 5.7 µg/m<sup>3</sup>, respectively)  
38  
39 205 using a breathing rate of 0.7 m<sup>3</sup>/h. Dermal uptake was then normalized by the air  
40  
41  
42 206 concentration during the 6 hours in the chamber and the participant’s weight. Also  
43  
44 207 reported is the average flux for the 6-h exposure, corrected for background uptake and  
45  
46 208 hood air inhalation. Exposed surface area is taken as 2.06 m<sup>2</sup>, estimated by equation 7A-7  
47  
48 209 of the Exposure Factors Handbook<sup>18</sup> and corrected for the area of the head (6.6% of total).  
49  
50  
51 210 To compare the rate of uptake among exposure conditions and between phthalates, we  
52  
53  
54 211 calculated a normalized metabolite excretion rate. First we calculated the slope of net  
55  
56 212 metabolite vs time from initial sample (after exposure begins) to the last sample that  
57  
58  
59  
60

includes no more than 75% of total net metabolite excreted. This slope divided by the total mass excreted was defined as the normalized metabolite excretion rate. See Table S.2 for additional details regarding the calculation methods.

## Results and Discussion

Major quantitative results are shown in Table 1. There is a striking difference between results for fresh and exposed clothing experiments. The net metabolites excreted over the 54 hour period after initiation of exposure are far higher for exposed clothes than for fresh clothes, indicating that parent compound uptake is much higher for exposed than for fresh clothes. When corrected for background uptake rate and inhalation from the breathing hood and normalized by body mass and air concentration, wearing exposed clothes resulted in DEP and DnBP uptakes that were 11 and 36 times greater, respectively, than when wearing fresh clothes. The mass of metabolites excreted over the first 24 hours by the volunteer wearing exposed clothes (3.6 mg MEP; 2.1 mg MnBP) approaches that due to application of a 2% DEP/DnBP cream over most of the skin of subjects as reported by Janjua et al.<sup>19</sup> (MEP range 2.5-85 mg; MnBP range 3.6-18 mg).

The ratio of the normalized uptake of DEP/DnBP was very different for the two scenarios. For exposed clothing, normalized uptake of DEP is somewhat smaller than for DnBP (DEP/DnBP = 0.7), but for fresh clothing it is much higher (DEP/DnBP = 2.3). This suggests that fresh clothes retard uptake of DnBP more than DEP and/or that exposed clothes enhance uptake of DnBP relative to DEP. Both mechanisms are consistent with a higher cloth/air partition coefficient for DnBP, which has a higher molecular weight than DEP.

1  
2  
3 236  
4  
5  
6 237 Normalized dermal uptake for both fresh and exposed clothes differ substantially from  
7  
8 238 uptake to bare skin as described in Weschler et al.<sup>8</sup> For comparison, results for bare skin  
9  
10 239 and clothing experiments are shown in Figure 1. For exposed clothes, normalized uptake of  
11  
12 240 DEP and DnBP are 3.3 and 6.5 times greater, respectively, than the average of bare skin  
13  
14 241 results and 1.9 and 3.9 times higher than the highest uptake observed in the bare skin  
15  
16 242 experiments. For fresh clothes, uptake is 3.2 and 5.6 times lower than the average for bare  
17  
18 243 skin experiments. Based on a t-test, the probability, *p*, of the results stemming from random  
19  
20 244 variation was  $<10^{-4}$  for exposed clothes and  $< 0.017$  for fresh clothes. These findings are  
21  
22 245 consistent with the hypothesis that fresh clothes retard uptake and exposed clothes  
23  
24 246 increase uptake compared with bare skin.  
25  
26  
27  
28  
29  
30 247  
31  
32 248 A comparison of the results, accounting for participant age, is also enlightening. Weschler  
33  
34 249 et al.<sup>8</sup> observed a striking relationship between dermal uptake and age. Shown in Figure 2a  
35  
36 250 and 2b are plots of net amounts of MEP and MnBP excreted, from the time exposure began  
37  
38 251 until the end of urine sampling, for the two clothing experiments (worn by a 48 year-old  
39  
40 252 participant) and bare skin results from the 47 year-old participant reported by Weschler et  
41  
42 253 al.<sup>8</sup> For both phthalates, wearing exposed clothing increased the excretion rate and net  
43  
44 254 excretion by a large margin. Wearing fresh clothes significantly reduced excretion rate and  
45  
46 255 net excretion.  
47  
48  
49  
50  
51 256  
52  
53  
54 257 In Figure 3, normalized clothing results for uptake of parent compounds are plotted against  
55  
56 258 age along with the normalized results for all six bare-skinned participants. The clothing  
57  
58  
59  
60

results are clearly “off the line”; the exposed clothes result in much higher uptake and fresh clothes much lower uptake than for bare skin. Again, using the 47 year-old participant from Weschler et al.<sup>8</sup> as the best comparator, we observe that exposed clothes resulted in 2.3 and 3.4 times more uptake of DEP and DnBP respectively. Fresh clothes resulted in 4.5 and 11 times lower uptake.

The excretion rate of metabolites differs among conditions and between phthalate metabolites. The difference between MnBP and MEP is apparent for exposed clothes in Figure 2, with MEP rising faster than MnBP. The normalized excretion rate for both conditions studied in this research and for the six bare skin participants is shown in Figure 4. To make the comparison more clear, the bare skin results for participants wearing hoods are grouped with the clothed results. Qualitatively, clothed results are similar to bare skin results: the normalized excretion rate of MEP is higher than for MnBP. For both MEP and MnBP, the excretion rate is higher for exposed clothes than for fresh clothes. This is consistent with the hypothesis that fresh clothes act as a barrier and delay transport from air to skin. The difference is more pronounced for MnBP than for MEP, possibly due to stronger sorption of DnBP to clothing.

The results support the hypotheses that 1) fresh clothes are protective, reducing uptake of DEP and DnBP compared with bare-skinned participants and 2) exposed clothes increase uptake. Although only one participant was tested (in two exposure periods), we believe the results are compelling, especially when compared with the narrow range of results for six bare-skinned participants. All results are significantly different from the six bare-skinned

1  
2  
3 282 participant results. When compared by age, the difference is even more apparent (Figure  
4  
5  
6 283 3). However, replication of these results with a larger number of participants will be  
7  
8 284 valuable.  
9  
10 285  
11  
12  
13 286 For both DEP and DnBP, the dose while wearing fresh clothes is small and could have come  
14  
15 287 from a combination of penetration through clothing and absorption by bare skin. The  
16  
17 288 participant in this study was not completely clothed: the hands were bare. We can estimate  
18  
19 289 the absorbed dose by hands assuming that hands are 4.7% of an average adult male's total  
20  
21 290 surface area.<sup>18</sup> We will use participant 2, the bare skinned participant closest in age to the  
22  
23 291 clothed participant, for comparison and will assume that the shorts worn by participant 2  
24  
25 292 covered approximately 5% of his total surface area. Correcting for the reduced total  
26  
27 293 exposed area due to hood (3.9% of total surface area) and shorts, the normalized dermal  
28  
29 294 uptake due to exposed hands for participant 2 would be approximately 0.004 and 0.003  
30  
31 295  $\mu\text{g}/\text{kg} / (\mu\text{g}/\text{m}^3)$  for DEP and DnBP respectively. These values can be compared with 0.017  
32  
33 296 and 0.007  $\mu\text{g}/\text{kg} / (\mu\text{g}/\text{m}^3)$  for DEP and DnBP for the fresh clothes experiment. Hence, for  
34  
35 297 DnBP, uptake by bare hands could represent a substantial fraction of total uptake from the  
36  
37 298 fresh clothing experiment. It is also interesting to note that the estimated DEP uptake by  
38  
39 299 bare hands accounts for only 25% of the observed uptake; therefore, penetration through  
40  
41 300 clothing may account for much of the uptake.  
42  
43 301  
44  
45 302 It is perhaps intuitive that fresh clothes should impede transfer from air to skin of airborne  
46  
47 303 contaminants. Clothing has been designed to protect workers from pesticide spray and  
48  
49 304 industrial toxic gases. A recent paper reported on the ability of "every-day" clothing<sup>20</sup> to  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 305 reduce in vitro dermal penetration of chlorpyrifos from solution. But early human subject  
4  
5  
6 306 studies of VOCs showed little influence of clothing on dermal absorption of nitrobenzene  
7  
8 307 and phenol.<sup>3,4</sup> This could be because sorption to cloth is weak for these low molecular  
9  
10 308 weight compounds. Higher molecular weight, low volatility organic compounds are known  
11  
12 309 to exhibit substantial air-fabric partitioning.<sup>21,22</sup> As volatility decreases, partitioning from  
13  
14 310 air to fabric increases and we would anticipate that retardation of transport across fabric  
15  
16 311 would also increase. Indeed, in this research we observed a lower normalized uptake of  
17  
18 312 DnBP relative to DEP, consistent with the roughly 25 times lower vapor pressure of DnBP.  
19  
20 313  
21  
22  
23  
24 314 Given the complicated geometry of fabric and skin, and the potential for air movement  
25  
26 315 through and under fabric, the data cannot be used to test more detailed models, to generate  
27  
28 316 exposure estimates or to identify compound/fabric combinations that would be most  
29  
30 317 protective or hazardous. Qualitatively, the transport of SVOCs into and out of clothing may  
31  
32 318 be described well by a model of transport of contaminants through porous media.<sup>23</sup> In the  
33  
34 319 context of this model, both advection and diffusion of contaminants through fabric would  
35  
36 320 be retarded by sorption. Key parameters influencing transport and dermal uptake are  
37  
38 321 likely to include the geometry and permeability of the fabric, how closely clothing fits, the  
39  
40 322 air-to-cloth partition coefficient, the dermal permeability of the contaminant, the elapsed  
41  
42 323 time the cloth is exposed to contaminated air after washing and the elapsed time clothing is  
43  
44 324 worn.  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 326 **Geometry and permeability of fabric.** Transport of air and moisture through fabric has  
55  
56 327 been extensively measured and modeled.<sup>24-27</sup> Hydraulic permeability lumps geometric  
57  
58  
59  
60



1  
2  
3 328 complexity of fabric into a parameter that characterizes fabric resistance to advective flux;  
4  
5  
6 329 hydraulic permeability is defined as the volume flux of air due to a specific pressure  
7  
8 330 difference across the fabric in units of  $\text{cm}^3/(\text{cm}^2 \text{ s})$  (usually at a pressure difference equal  
9  
10 331 to 125 Pa). Hydraulic permeability can range over several orders of magnitude: very low  
11  
12 332 ( $<0.1 \text{ cm}^3/(\text{cm}^2 \text{ s})$ ) for dense or sealed materials and very high for loosely woven thin  
13  
14 333 fabrics ( $>300 \text{ cm}^3/(\text{cm}^2 \text{ s})$ ). SVOC transport is likely to be more influenced by advection in  
15  
16 334 loosely woven materials with a high hydraulic permeability; for tightly woven materials,  
17  
18 335 diffusive transport is expected to dominate. For intermediate materials, the relative  
19  
20 336 contributions of diffusion and advection will be influenced by pressure gradients, wind and  
21  
22 337 movement.  
23  
24  
25  
26 338

27  
28  
29 339 **How close clothing fits.** Some sorbed SVOCs may transfer from cloth to skin by contact.  
30  
31 340 However, since most of the surface area available for adsorption in a woven fabric is  
32  
33 341 internal, only a small fraction of the sorbed SVOC is likely a consequence of transfer by  
34  
35 342 direct contact with the outer fabric fibers. Instead, we believe that the more important  
36  
37 343 mechanism is desorption from fiber surfaces and diffusion across a thin air gap to skin. For  
38  
39 344 diffusion across a quiescent air gap, flux is proportional to the reciprocal of the air gap  
40  
41 345 distance. The air gap distance was not measured in this study but we estimate it ranged  
42  
43 346 from  $<0.1 \text{ cm}$  to  $0.5 \text{ cm}$ . By comparison, a typical bare-skin concentration boundary layer is  
44  
45 347 about  $0.2$  to  $0.4 \text{ cm}$ , which can be estimated by dividing the gas diffusivity of the SVOC  
46  
47 348 ( $0.056 \text{ cm}^2/\text{s}$ )<sup>28</sup> by an air-to-skin deposition velocity ( $0.14$ - $0.28 \text{ cm/s}$ )<sup>29</sup>. Therefore, the  
48  
49 349 initial flux from fabric to skin could be smaller, or more than 4 times greater, than from  
50  
51 350 bulk air when wearing equilibrated clothing. Notably, this estimate overlaps the observed  
52  
53  
54  
55  
56  
57  
58  
59  
60

ratio of uptake for exposed clothing to the average for bare skin (3.3 for DEP and 6.5 for DnBP).

**Air-to-cloth partitioning.** The sorptive capacity of fabric will influence how it reduces or enhances transport from air to skin. There is recognition that adsorption and desorption of indoor-relevant gases on fabrics for tobacco smoke products<sup>30-32</sup> and pesticides<sup>21</sup> can influence exposure. Several studies have shown that dry-cleaning solvents<sup>33-36</sup> and moth repellants<sup>37</sup> can sorb to clothing and subsequently desorb, increasing indoor concentrations. Specialty fabrics have been developed that sorb or react with chemical warfare agents or pesticides to protect the wearer.<sup>38</sup> However, we have only identified two papers<sup>22,39</sup> that report equilibrium partition coefficients for an indoor air contaminant and commonly worn fabrics. In one paper<sup>22</sup> the investigators measured equilibrium partition coefficients for airborne free-base methamphetamine and fabrics including cotton and polyester. The partition coefficients were high enough that mouthing of these fabrics was predicted to be the primary route of exposure for toddlers, similar to the observation by Gurunathan et al.<sup>21</sup> for chlorpyrifos and plush toys.

**Dermal permeability.** We anticipate that the compounds that are most likely to exhibit enhanced dermal uptake from exposed clothes are those that have high dermal permeability coupled with gas-to-fabric partition coefficients in an intermediate range (not too high, not too low). Uptake of compounds with low dermal permeability is limited by resistance across skin; modest changes in mass-transfer conditions external to skin will likely have little impact on overall uptake. Dermal permeability of compounds typical of

1  
2  
3 374 indoor air have been estimated by Weschler and Nazaroff.<sup>6</sup> They identified more than 30  
4  
5  
6 375 common indoor pollutants that are predicted to have high dermal uptake relative to  
7  
8 376 inhalation uptake. If a compound has too high a gas-to-fabric partition coefficient, this will  
9  
10 377 retard transfer from the fabric to skin. On the other hand, if a compound has too small a  
11  
12 378 gas-to-fabric partition coefficient, then exposed clothes have sorbed very little of the  
13  
14 379 compound and there will be concomitantly little enhancement. It is in the intermediate  
15  
16 380 range of gas-fabric partitioning that sorption to clothes prior to wear will have the greatest  
17  
18 381 enhancement on uptake.  
19

20 382  
21  
22  
23  
24  
25 383 **Elapsed time clothing sorbs contaminants and time clothing is worn.** It takes time for  
26  
27 384 fabric to adsorb airborne contaminants and approach equilibrium with gas phase  
28  
29 385 concentrations. It also takes time for contaminants to desorb and transfer to skin. As an  
30  
31 386 example, consider a tight-fitting shirt that has been washed, stored in the presence of a  
32  
33 387 contaminant (in air) and then worn. If we assume that the characteristic time for the fabric  
34  
35 388 to equilibrate ( $\tau_e$ ) is independent of the air concentration, then we can qualitatively  
36  
37 389 compare this time to the actual time stored in the presence of a contaminant ( $t_s$ ) or the time  
38  
39 390 clothing is worn after storage ( $t_w$ ). *Scenario 1:*  $t_s < \tau_e$ . For this scenario, the fabric has not  
40  
41 391 equilibrated with the contaminant concentration in the air and may in fact continue to sorb  
42  
43 392 contaminants even while worn. Regardless of the chemical, enhanced flux from cloth to  
44  
45 393 skin will be limited. *Scenario 2:*  $t_s \geq \tau_e$  and  $t_w < \tau_e$ . For this scenario the fabric is well  
46  
47 394 equilibrated with a contaminant before wearing, but the time worn is short relative to the  
48  
49 395 time it takes for the contaminant to reach a new steady-state. During the time worn, flux to  
50  
51 396 skin will be enhanced but the mass adsorbed to fabric will not change substantially. This  
52  
53  
54  
55  
56  
57  
58  
59  
60

scenario could be represented by a shirt worn for a short time that had adsorbed a relatively more volatile, low partition coefficient chemical; it could also be represented by a shirt worn for a longer period of time that had adsorbed a less volatile, higher partition coefficient chemical. *Scenario 3*:  $t_s \geq \tau_e$  and  $t_w > \tau_e$ . Here, the shirt has been worn long enough that a substantial fraction of the contaminant has desorbed. While the initial flux to skin may be high, the time-averaged flux will be lower than for *Scenario 2* (all else being equal).

Since DnBP is anticipated to have a higher partition coefficient than DEP, we would anticipate that  $\tau_e$  would be greater for DnBP than DEP. We observe a normalized dermal uptake that is higher for DnBP than DEP from exposed clothes. If the 6-hour period that the participant wears the exposed clothing is similar or longer than  $\tau_e$  for DEP, then it may fall under *Scenario 3*, while DnBP falls under *Scenario 2*.

## Conclusions

Clothing acts as a barrier to exposure, but also as a reservoir for recently adsorbed chemicals; the latter can increase dermal uptake. Not only are people subjected to airborne SVOCs while at home, they are also exposed to “home pollutants” outside of their residence when they wear clothing that has been stored in the presence of various SVOCs at home. Given the very large increase in the normalized dermal uptake of DEP and DnBP observed for exposed fabric in this study, we believe clothing-mediated dermal uptake is an under-recognized exposure pathway that could be a substantial or even a dominant exposure route for many chemicals. This is of potential importance in occupational as well as non-occupational settings.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

420

421   **Acknowledgements**

422   The authors are grateful to Louise B. Weschler for her assistance with these experiments.

423

424

For Peer Review Only

## References

- 1 Rehal B, Maibach H. Percutaneous absorption of vapors in human skin. *Cutan Ocul Toxicol* 2011; **30**: 87–91.
- 2 Rauma M, Boman A, Johanson G. Predicting the absorption of chemical vapours. *Adv Drug Deliv Rev* 2013; **65**: 306–314.
- 3 Piotrowski J. Further investigations on the evaluation of exposure to nitrobenzene. *Br J Ind Med* 1967; **24**: 60–65.
- 4 Piotrowski J. Evaluation of exposure to phenol: absorption of phenol vapour in the lungs and through the skin and excretion of phenol in urine. *Br J Ind Med* 1971; **28**: 172–178.
- 5 Weschler CJ, Nazaroff WW. SVOC exposure indoors: Fresh look at dermal pathways. *Indoor Air* 2012; **22**: 356–377.
- 6 Weschler CJ, Nazaroff WW. Dermal uptake of organic vapors commonly found in indoor air. *Environ Sci Technol* 2014; **48**: 1230–1237.
- 7 Gong M, Zhang Y, Weschler CJ. Predicting dermal absorption of gas-phase chemicals: Transient model development, evaluation, and application. *Indoor Air* 2014; **24**: 292–306.
- 8 Weschler CJ, Beko G, Koch HM, Salthammer T, Schripp T, Toftum J *et al.* Transdermal uptake of diethyl- and di(n-butyl) phthalate directly from air: experimental verification. *Environ Health Perspect* 2015; doi:10.1289/ehp.1409151.
- 9 Blum A, Gold MD, Ames BN, Jones FR, Hett EA, Dougherty RC *et al.* Children absorb tris-BP flame retardant from sleepwear: urine contains the mutagenic metabolite, 2,3-dibromopropanol. *Science* 1978; **201**: 1020–1023.
- 10 Appel KE, Gundert-Remy U, Fischer H, Faulde M, Mross KG, Letzel S *et al.* Risk assessment of Bundeswehr (German Federal Armed Forces) permethrin-impregnated battle dress uniforms (BDU). *Int J Hyg Environ Health* 2008; **211**: 88–104.
- 11 Rossbach B, Appel KE, Mross KG, Letzel S. Uptake of permethrin from impregnated clothing. *Toxicol Lett* 2010; **192**: 50–55.
- 12 Kegel P, Letzel S, Rossbach B. Biomonitoring in wearers of permethrin impregnated battle dress uniforms in Afghanistan and Germany. *Occup Environ Med* 2014; **71**: 112–117.

- 1  
2  
3 458 13 Schripp T, Salthammer T, Fauck C, Beko" G, Weschler CJ. Latex paint as a delivery  
4 459 vehicle for diethylphthalate and di-n-butylphthalate: Predictable boundary layer  
5 460 concentrations and emission rates. *Sci Total Environ* 2014; **494**: 299–305.  
6  
7  
8 461 14 Koch HM, Gonzalez-Reche LM, Angerer J. On-line clean-up by multidimensional liquid  
9 462 chromatography-electrospray ionization tandem mass spectrometry for high  
10 463 throughput quantification of primary and secondary phthalate metabolites in human  
11 464 urine. *J Chromatogr B Analyt Technol Biomed Life Sci* 2003; **784**: 169–182.  
12  
13  
14 465 15 Koch HM, Christensen KLY, Harth V, Lorber M, Brüning T. Di-n-butyl phthalate (DnBP)  
15 466 and diisobutyl phthalate (DiBP) metabolism in a human volunteer after single oral  
16 467 doses. *Arch Toxicol* 2012; **86**: 1829–1839.  
17  
18  
19 468 16 Koch HM, Becker K, Wittassek M, Seiwert M, Angerer J, Kolossa-Gehring M. Di-n-  
20 469 butylphthalate and butylbenzylphthalate - Urinary metabolite levels and estimated  
21 470 daily intakes: Pilot study for the German Environmental Survey on children. *J Expo Sci*  
22 471 *Environ Epidemiol* 2007; **17**: 378–387.  
23  
24  
25 472 17 Koch HM, Lorber M, Christensen KLY, Pälme C, Koslitz S, Brüning T. Identifying  
26 473 sources of phthalate exposure with human biomonitoring: Results of a 48h fasting  
27 474 study with urine collection and personal activity patterns. *Int J Hyg Environ Health*  
28 475 2013; **216**: 672–681.  
29  
30  
31 476 18 USEPA. Exposure Factors Handbook: 2011 Edition. US Environmental Protection  
32 477 Agency: Washington D.C, 2011.  
33  
34 478 19 Janjua NR, Frederiksen H, Skakkebaek NE, Wulf HC, Andersson A-M. Urinary excretion  
35 479 of phthalates and paraben after repeated whole-body topical application in humans. *Int*  
36 480 *J Androl* 2008; **31**: 118–129.  
37  
38  
39 481 20 Moya J, Howard-Reed C, Corsi RL. Volatilization of chemicals from tap water to indoor  
40 482 air from contaminated water used for showering. *Environ Sci Technol* 1999; **33**: 2321–  
41 483 2327.  
42  
43 484 21 Moore CA, Wilkinson SC, Blain PG, Dunn M, Aust GA, Williams FM. Use of a human skin  
44 485 in vitro model to investigate the influence of 'every-day' clothing and skin surface  
45 486 decontamination on the percutaneous penetration of organophosphates. *Toxicol Lett*  
46 487 2014; **229**: 257–264.  
48  
49 488 22 Gurunathan S, Robson M, Freeman N, Buckley B, Roy A, Meyer R *et al*. Accumulation of  
50 489 chlorpyrifos on residential surfaces and toys accessible to children. *Environ Health*  
51 490 *Perspect* 1998; **106**: 9–16.  
52  
53  
54 491 23 Morrison GC, Shakila N, Parker K. Accumulation of gas-phase methamphetamine on  
55 492 clothing , toy fabrics and skin oil. *Indoor Air* 2014; doi: 10.1111/ina.12159.  
56  
57  
58  
59  
60



- 493 24 Fetter CW. *Contaminant Hydrogeology*. MacMillan Publishing Company: New York,  
494 1993.
- 495 25 Mavruz S, Ogulata RT. Investigation of air permeability of single jersey fabrics with  
496 different relaxation states. *J Text Inst* 2011; **102**: 57–64.
- 497 26 Ogulata RT, Mavruz S. Investigation of porosity and air permeability values of plain  
498 knitted fabrics. *Fibres Text East Eur* 2010; **82**: 71–75.
- 499 27 Gibson WB, Keller PR, Foltz DJ, Harvey GJ. Diethylene glycol mono butyl ether  
500 concentrations in room air from application of cleaner formulations to hard surfaces. *J*  
501 *Expo Anal Environ Epidemiol* 1991; **1**: 369–383.
- 502 28 Ghaddar N, Ghali K, Harathani J, Jaroudi E. Ventilation rates of micro-climate air  
503 annulus of the clothing-skin system under periodic motion. *Int J Heat Mass Transf* 2005;  
504 **48**: 3151–3166.
- 505 29 Lyman W, Reehl W, Rosenblatt D. *Handbook of Chemical Property Estimation Methods*.  
506 Ameican Chemical Society, 1990.
- 507 30 Weschler CJ, Salthammer T, Fromme H. Partitioning of phthalates among the gas phase,  
508 airborne particles and settled dust in indoor environments. *Atmos Environ* 2008; **42**:  
509 1449–1460.
- 510 31 Noble RE. Environmental tobacco smoke uptake by clothing fabrics. *Sci Total Environ*  
511 2000; **262**: 1–3.
- 512 32 Chien Y-C, Chang C-P, Liu Z-Z. Volatile organics off-gassed among tobacco-exposed  
513 clothing fabrics. *J Hazard Mater* 2011; **193**: 139–148.
- 514 33 Ueta I, Saito Y, Teraoka K, Miura T, Jinno K. Determination of volatile organic  
515 compounds for a systematic evaluation of third-hand smoking. *Anal Sci* 2010; **26**: 569–  
516 574.
- 517 34 Tichenor BA, Sparks LE, Jackson MD, Guo Z, Mason Plunket MACM, Rasor SA. Emissions  
518 of perchloroethylene from dry cleaned fabrics. *Atmospheric Environ - Part Gen Top*  
519 1990; **24 A**: 1219–1229.
- 520 35 Chao CYH, Tung TCW, Niu JL, Pang SW, Lee RYM. Indoor perchloroethylene  
521 accumulation from dry cleaned clothing on residential premises. *Build Environ* 1998;  
522 **34**: 319–328.
- 523 36 Thomas KW, Pellizzari ED, Perritt RL, Nelson WC. Effect of dry-cleaned clothes on  
524 tetrachloroethylene levels in indoor air, personal air, and breath for residents of several  
525 New Jersey homes. *J Expo Anal Environ Epidemiol* 1991; **1**: 475–490.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

526 37 Sherlach KS, Gorka AP, Dantzler A, Roepe PD. Quantification of perchloroethylene  
527 residues in dry-cleaned fabrics. *Environ Toxicol Chem* 2011; **30**: 2481–2487.

528 38 Guerrero PA, Corsi RL. Emissions of p-dichlorobenzene and naphthalene from  
529 consumer products. *J Air Waste Manag Assoc* 2012; **62**: 1075–1084.

530 39 Schreuder-Gibson H, Truong Q, Walker J, Owens J, Wander J, Jones W. Chemical and  
531 Biological Protection and Detection in Fabrics for Protective Clothing. *MRS Bull* 2003;  
532 **28**: 574–578.

533 40 Piadé JJ, D’Andrés S, Sanders EB. Sorption phenomena of nicotine and ethenylpyridine  
534 vapors on different materials in a test chamber. *Environ Sci Technol* 1999; **33**: 2046–  
535 2052.

536

**Table legend**

**Table 1.** Net amount of metabolites excreted during the time period from entering the chamber until 54 hours later, as well as parent compound uptake calculated from the metabolite levels; details regarding the calculation of the listed values are presented in Supporting Information.

For Peer Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure legends.**

**Figure 1.** Normalized mass of DEP and DnBP absorbed for fresh and exposed clothes experiments. Also shown for comparison are results from the 6 bare-skinned participants (boxplot) reported in Weschler et al.<sup>8</sup> The line within the box represents the median; the bottom and top of the box, the 25<sup>th</sup> and 75<sup>th</sup> percentiles; the upper and lower whiskers, the 10<sup>th</sup> and 90<sup>th</sup> percentiles.

**Figure 2.** Net amount of MEP (2a) and MnBP (2b) excreted from beginning of exposure until last urine sample. Results for fresh and exposed clothes are compared against the bare skin results of the closest aged participant in Weschler et al.<sup>8</sup>

**Figure 3.** Normalized dermal uptake of DEP and DnBP versus age. Shown are results from this research (clothes) and results for six bare skin participants reported by Weschler et al.<sup>8</sup>

**Figure 4.** Normalized metabolite excretion rate for MEP and MnBP. Shown are results from this research (clothes) and results for six bare-skinned participants reported by Weschler et al.<sup>8</sup> The line within the box represents the median; the bottom and top of the box, the 25<sup>th</sup> and 75<sup>th</sup> percentiles; the upper and lower whiskers, the 10<sup>th</sup> and 90<sup>th</sup> percentiles.




Figure 1. Normalized mass of DEP and DnBP absorbed for fresh and exposed clothes experiments. Also shown for comparison are results from the 6 bare-skinned subjects (boxplot) reported in Weschler et al.<sup>8</sup> The line within the box represents the median; the bottom and top of the box, the 25th and 75th percentiles; the upper and lower whiskers, the 10th and 90th percentiles.

114x80mm (300 x 300 DPI)




Figure 3. Normalized dermal uptake of DEP and DnBP versus age. Shown are results from this research (clothes) and results for six bare skin subjects reported by Weschler et al.<sup>8</sup>  
241x175mm (72 x 72 DPI)





Figure 2. Net amount of MEP (2a) and MnBP (2b) excreted from beginning of exposure until last urine sample. Results for fresh and exposed clothes are compared against the bare skin results of the closest aged subject in Weschler et al.<sup>8</sup>  
121x50mm (300 x 300 DPI)



Normalized metabolite excretion rate for MEP and MnBP. Shown are results from this research (clothes) and results for six bare-skinned subjects reported by Weschler et al.<sup>8</sup> The line within the box represents the median; the bottom and top of the box, the 25th and 75th percentiles; the upper and lower whiskers, the 10th and 90th percentiles.  
129x108mm (300 x 300 DPI)

**Table 1.** Net amount of metabolites excreted during the time period from entering the chamber until 54 hours later, as well as parent compound uptake calculated from the metabolite levels; details regarding the calculation of the listed values are presented in Supporting Information.

	Metabolites excreted (µg)			Total uptake parent (µg)		Background corrected uptake parent (µg)		Dermal only uptake parent (corrected for concentration in hood) (µg)		Normalized dermal uptake (µg/kg/(µg/m <sup>3</sup> ))		Average flux based on 6 hour exposure (µg /m <sup>2</sup> /h)	
	MEP	MnBP	3OH-MnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP
<b>Fresh clothing</b>	466	121	7.7	634	176	522	98	352	74	0.017	0.007	28	6
<b>Exposed clothing</b>	3666	2367	136	4995	3432	4882	3355	4712	3331	0.178	0.261	381	270



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Supplemental Material**

**Role of clothing in both increasing and decreasing dermal absorption of airborne SVOCs**

Glenn Morrison<sup>1</sup>, Charles J. Weschler<sup>2,3</sup>, Gabriel Bekö<sup>2</sup>, Holger Koch<sup>4</sup>, Tunga Salthammer<sup>5</sup>, Tobias Schripp<sup>5</sup>, Jørn Toftum<sup>2</sup> and Geo Clausen<sup>2</sup>

<sup>1</sup>Civil, Architectural and Environmental Engineering, Missouri University of Science and Technology, 1401 N. Pine St., Rolla, MO 65409, USA

<sup>2</sup>International Centre for Indoor Environment and Energy, Department of Civil Engineering, Technical University of Denmark, DK-2800 Lyngby, Denmark

<sup>3</sup>Environmental and Occupational Health Sciences Institute, Rutgers University, 170 Frelinghuysen Road, Piscataway, NJ 08854, USA

<sup>4</sup>Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr-University Bochum (IPA), Bürkle-de-la-Camp-Platz 1, 44789 Bochum, Germany

<sup>5</sup>Fraunhofer WKI, Department of Material Analysis and Indoor Chemistry, Bienroder Weg 54E, 38108 Braunschweig, Germany

# Table of Contents for Supplementary Materials

## Page

Table S1. Clothing specifications determined from product packaging or labels 3

Figure S1. Male subject shown wearing full set of test clothing and the breathing hood while seated in the test chamber 3

Table S2. Metabolites excreted and parent compound uptake 4

For Peer Review Only

**Table S1.** Clothing characteristics determined from product packaging or labels

Clothing	Composition	Size	Manufacturer	Description
Undershirt	100% cotton	M/M, 38-40" (97-102 cm)	Gildan	Short sleeve, crew neck, color: white Estimated cloth area = 0.91 m <sup>2</sup>
Underwear	100% cotton with elastic band	L/G, 36-38" (91-97 cm)	Hanes	Boxer style briefs, color: grey Estimated cloth area = 0.24 m <sup>3</sup>
Shirt	100% cotton	M	Gildan	Long sleeve tee-shirt, crew neck, color: dark green Estimated cloth area = 1.03 m <sup>3</sup>
Pants	100% cotton	36" (91 cm) waist 36" (91 cm) inseam	Wrangler	Jeans, slim fit, color: dark blue Estimated cloth area = 1.10 m <sup>2</sup>
Socks	85% cotton 12% polyester 1% elastic 1% nylon 1% spandex	12W-15	Starter	Tube socks that rise ~20 cm above ankle, color: white Estimated cloth area (pair) = 0.07 m <sup>2</sup>



**Figure S1.** Male subject shown wearing full set of test clothing and the breathing hood while seated in the test chamber.

**Table S2.** Net amount of metabolites excreted during the time period from entering the chamber until 54 hours later, as well as parent compound uptake calculated from the metabolite levels. This Table is identical to Table 1, but is included here with detailed explanations of calculation methods.

	Metabolites excreted ( $\mu\text{g}$ ) <sup>1</sup>			Total uptake parent ( $\mu\text{g}$ ) <sup>2</sup>		Background corrected uptake parent ( $\mu\text{g}$ ) <sup>3</sup>		Dermal only uptake parent (corrected for concentration in hood) ( $\mu\text{g}$ ) <sup>4</sup>		Normalized dermal uptake ( $\mu\text{g}/\text{kg}/(\mu\text{g}/\text{m}^3)^5$		Average flux ( $\mu\text{g}/\text{m}^2/\text{h}$ ) <sup>6</sup>	
	MEP	MnBP	3OH-MnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP	DEP	DnBP
Fresh clothing	466	121	7.7	634	176	522	98	352	74	0.017	0.007	28	6
Exposed clothing	3666	2367	136	4995	3432	4882	3355	4712	3331	0.178	0.261	381	270

1. The mass of metabolites excreted is determined by multiplying the concentration of each metabolite by the volume of urine collected for each sample and summing over all samples collected during the 54 hour period after the exposure started.

2. Total parent uptake is calculated by converting mass from metabolite to parent and using a metabolic conversion factor.

Compound	Abbreviation	CAS-no.	Molecular weight (g/mol)	Metabolic conversion factor
Diethylphthalate	DEP	84-66-2	222.24	NA
Di-n-butylphthalate	DnBP	84-74-2	278.34	NA
Monoethylphthalate	MEP	2306-33-4	194.18	0.84
Mono-n-butylphthalate	MnBP	131-70-4	222.24	0.84
3OH-mono-n-butylphthalate	3OH-MnBP	57074-43-8	238.24	0.07

$$\text{DEP} = [(\text{MEP} / 194.18) * 222.24] / 0.84$$

$$\text{DnBP} = [(\text{MnBP} / 222.24) * 278.34] + [(3\text{OH-MnBP} / 238.24) * 278.34] / (0.84 + 0.07)$$

3. Background-corrected uptake of the parent compound is determined by subtracting out the background concentration of metabolites, integrating the resulting mass, then applying the conversion described in (2) above. Background is defined as the pre-exposure urine concentration.

4. Dermal uptake of parent compounds is calculated by subtracting from background-corrected uptake the inhaled mass of DEP and DnBP based on concentrations in breathing air of the hood ( $40.7$  and  $5.7 \mu\text{g}/\text{m}^3$ , respectively). Inhalation rate is assumed to be  $0.7 \text{ m}^3/\text{h}$ . Therefore, the mass subtracted is  $170$  and  $24 \mu\text{g}$  for DEP and DnBP respectively.

5. Normalized uptake is calculated by dividing the dermal uptake by average exposure air concentration and the subject body mass. Average air concentrations during the fresh clothing experiment were  $230 \mu\text{g}/\text{m}^3$  DEP and  $113$

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

$\mu\text{g}/\text{m}^3$  DnBP. Average air concentrations during the exposed clothing experiment were 291  $\mu\text{g}/\text{m}^3$  DEP and 140  $\mu\text{g}/\text{m}^3$  DnBP.

6. The average flux is estimated from the “Dermal only” corrected parent compound uptake, divided by exposed surface area of the participant and the exposure period (6 hours). Exposed surface area is taken as 2.06  $\text{m}^2$ , estimated by equation 7A-7 of the Exposure Factors Handbook<sup>18</sup> and corrected for the area of the head (6.6% of total).

For Peer Review Only